PREMENSTRUAL SYMPTOM TRACKER

(DAILY RECORD OF SEVERITY OF PROBLEMS)

Name: Month:



INSTRUCTIONS

Print off as many copies as you need to complete a full two months worth of tracking. Begin tracking your premenstrual symptoms with this chart today. Fill it out daily (preferably at the end of your day). Two full months of menstrual cycle charting will allow for a more accurate assessment.

Each evening note the degree to which you experienced each of the problems listed below. Put an "x" in the box which corresponds to the severity:

Enter day of the week (e.g. Monday =	'M')																															
Note any spotting by entering																															_	
Note menstrual bleeding by entering	g 'M'		Ш																												L	
Date (i.e. 1 = 1st of the mo	nth)	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	3
Felt depressed, sad, "down,", or	6 5		-																													\vdash
"blue" or felt hopeless; or felt	4																															
worthless or guilty	3																														_	
	2																														\vdash	\vdash
2. Felt anxious, tense, "keyed up" or	6																															
"on edge"	5 4																-														\vdash	+
	3																															t
	2																															I
	6		\vdash																												\vdash	┢
3. Had mood swings (i.e., suddenly	5																															
feeling sad or tearful) or was	4																															\perp
sensitive to rejection or feelings	2		\vdash																												\vdash	+
were easily hurt	1																															
4. Felt angry, or irritable	6																															$oxed{\Box}$
To ron angry, or innable	5 4		\vdash																												\vdash	+
	3																															_
	2		\prod																												<u> </u>	+
- 11	6																															+
5. Had less interest in usual activities	5																															_
(work, school, friends, hobbies)	4		\prod																												<u> </u>	\perp
	2		\vdash																													+
	1																															
6. Had difficulty concentrating	6		\square																												\vdash	\perp
	5 4		\vdash																												\vdash	H
	3																															
	2																														\vdash	_
	6																															╁
7. Felt lethargic, tired, or fatigued; or had lack of energy	5																															
	4																														<u> </u>	
	2																															H
	1																															
8. Had increased appetite or overate; or had cravings for specific foods	6 5																															
	4																															H
	3																															
	2																															\vdash
	6																															T
9. Slept more, took naps, found it hard																																
to get up when intended; or had	3																															\vdash
trouble getting to sleep or staying asleep	2																															
usieep	1																															L
10. Felt overwhelmed or unable to cope;	6 5																														\vdash	\vdash
or felt out of control	4																															
	3																														<u> </u>	L
	2																														\vdash	\vdash
11. Had breast tenderness, breast	6																															
swelling, bloated sensation, weight	5																										\Box		\dashv		\vdash	
gain, headache, joint or muscle	3																\dashv				-						\dashv		\dashv		\vdash	H
pain, or other physical symptoms	2																															
	1																															Ļ
At work, school, home, or in daily	6 5																												\dashv		\vdash	
routine, at least one of the problems	4																															
noted above caused reduction of	2																														\vdash	-
production of efficiency	1																															-
And the second s	6																															
At least one of the problems noted above caused avoidance of or less	5 4														<u> </u>														-		\vdash	-
participation in hobbies or social	3																															F
activities	2																															
	1																														 	-
At least one of the problems noted	6 5																															
above interfered with relationships	4																															
with others	2														<u> </u>		\vdash														<u> </u>	-
															I '										'		'				1	

PREMENSTRUAL SYMPTOM TRACKER

(DAILY RECORD OF SEVERITY OF PROBLEMS)

Name:

Month:

INSTRUCTIONS

Print off as many copies as you need to complete full two months worth of tracking. Begin tracking your premenstrual symptoms with this chart today. Fill it out daily (preferably at the end of your day)

Each evening note the dear blems listed below. Put an "x" in the box which corresponds to the severity:

1 - not at all **2 - minim** and **4 - moderate** 5 - seve strem

ı – no	ot at all 2 - minir	4 -	mo	aero	ale	J .	- se	VC			AIL	eme	• 																				
	Enter day of th g. Monday = '	'M')	S	S	m	†	W	†	f	N																							
	Not ting by entering																																
	Note mer eding by entering	'M'																		Ц													
	i.e. 1 = 1st of the mor	nth)	1	2	3	4	5	6	7	8	9			12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31
1.	Felt deprad, "down,", or	6 5									\ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \									\dashv													
	"blue" or peless; or felt	4									X	1																					
	worthles ty	2		X				X	X	X		+								\dashv		\dashv								\vdash			
		1	X	^	Χ	Χ	Χ													_													
2.	Felt anxiesse, "keyed up" or	6 5														-				\dashv		\dashv											
	"on edge	4									X																						
		2	X	X		X	X	X	X	X										\dashv		-											
_		1			Χ						7									_												\Box	
3.	Had mood swii uddenly	6 5		_																-													
	feeling sad or tecanons	4							X																								
	sensitive to rejection	2			X	X	У	X												-													
	were easily hurt	2		V		**																										\exists	
4.	Felt angry, or irritable																																
		4																															
	-	2		_																V						_				\square			
		1																															
5.	Had less interest in usual activities	6 5																															
	(work, school, friends, hobbies)	4																					1										
		2		_																			}										
		1																															
6	Had difficulty concentrating	6 5		_																													
0.		4																												\square			
		3																		-												\Box	
		1																		\dashv													
7	Felt lethargic, tired, or fatigued; or	6 5																				-								\Box		\neg	
/ •	had lack of energy	4																															
	O ,	3																		-												$\overline{}$	
		1																															
8.	Had increased appetite or overate;	6 5																												\Box			
•	or had cravings for specific foods	4																															
		2		_																_		-								\square			
		1																															
9.	Slept more, took naps, found it hard	6 5																		-										\square			
	to get up when intended; or had	4																															
	trouble getting to sleep or staying	2																		\dashv										$\vdash\vdash$		\rightarrow	
	asleep	1																												\square		\rightrightarrows	
10	Felt overwhelmed or unable to cope;	6 5																												$\vdash\vdash$		\rightarrow	
	or felt out of control	4																												口		\exists	
		2																		\dashv		_								$\vdash\vdash$		\rightarrow	
		1																														\rightrightarrows	
11.	Had breast tenderness, breast	6 5														\dashv				\dashv		\dashv								$\vdash\vdash$		\dashv	
	swelling, bloated sensation, weight	4																												口			
	gain, headache, joint or muscle pain, or other physical symptoms	2																		\dashv		\dashv								$\vdash\vdash$		\dashv	
	pain, or other physical symptoms	1																		_												\exists	
	At work, school, home, or in daily	6 5																												$\vdash \vdash$			
	routine, at least one of the problems	4																												口		\Box	
	noted above caused reduction of	2																		\dashv		\dashv								$\vdash \vdash$		\dashv	
	production of efficiency	-1																												口		\Box	
	At least one of the problems noted	6 5																		+		\dashv								$\vdash \vdash$		\dashv	
	above caused avoidance of or less	4																												口		\Box	
l	participation in hobbies or social	2																				\dashv								$\vdash \vdash$		\dashv	
1	activities	1																														\rightrightarrows	
] .	At least one of the problems noted	6 5																				\dashv								$\vdash \vdash$		\rightarrow	
	above interfered with relationships	4																												\Box		\dashv	
1	with others	2																		\dashv		\dashv						_		$\vdash \vdash$		\dashv	
		-												 		$\overline{}$		-		_		-+								$\overline{}$		\rightarrow	